



<http://www.biodiversitylibrary.org/>

**The Journal of the Bombay Natural History Society.**

Bombay :The Society,

<http://www.biodiversitylibrary.org/bibliography/7414>

**v.100 (2003):** <http://www.biodiversitylibrary.org/item/189679>

Page(s): Page 631

Contributed by: Smithsonian Libraries

Sponsored by: Biodiversity Heritage Library

Generated 20 January 2016 2:00 AM

<http://www.biodiversitylibrary.org/pdf4/047527500189679>

This page intentionally left blank.

We are grateful to Mr. B.C. Choudhury, Nodal Officer, GOI-UNDP Sea Turtle Project, and Dr. V.S. Vijayan, Director, SACON, for encouraging us throughout this study. Our colleagues P.R. Arun and M. Rajasekaran helped us in preparing the map.

March 13, 2002

S. BHUPATHY  
S. SARAVANAN  
Sálim Ali Centre for Ornithology  
and Natural History,  
P.O. Anaikatti, Coimbatore 641 108,  
Tamil Nadu, India.

## REFERENCES

- AGASTHEESAPILLAI, A. & R. THIAGARAJAN (1979): Biology of the green turtle *Chelonia mydas* (Linnaeus) in the Gulf of Mannar and Palk Bay. *J. mar. biol. Assn India* 21(1&2): 45-60.
- DAS, I. (1995): Turtles and Tortoises of India. Oxford University Press, Bombay. 176 pp.
- FRAZIER, J. (1980): Exploitation of marine turtles in the Indian Ocean. *Human Ecol.* 8(4): 329-370.
- KAR, C.S. & S. BHASKAR (1995): Status of sea turtles in the Indian Ocean. Pp. 356-372. *In: Biology and conservation of sea turtles* (Ed.: Bjorndal, K.A.). Smithsonian Institution Press, Washington, DC.
- KURIYAN, G.K. (1950): Turtle fishing in the sea around Krusadai Island. *J. Bombay nat. Hist. Soc.* 49: 509-512.

9. RECORD OF THE PAINTED KALOULA *KALOULA TAPROBANICA* IN ANDHRA PRADESH

The painted kaloula, *Kaloula taprobanica* Parker, is a medium sized (5.0-8.0 cm) frog that prefers urban areas and open forests (Daniels 1997). The species is reported from Assam, West Bengal, Bihar, Madhya Pradesh, Orissa, Karnataka and Tamil Nadu (Dutta 1997). The species has not been reported from Andhra Pradesh, though reported from its four bordering states (Orissa and Madhya Pradesh in the north, Karnataka to the west and Tamil Nadu to the south) (Daniel 2002, Dutta 1997).

During our studies on the herpetofauna in Sriharikota Island, Nellore district, Andhra Pradesh, we found the species to be one of the most common frogs on the Island. Sriharikota borders Tamil Nadu, and records of the species in this State are from its eastern areas (Daniels 1997). Specimens have also been collected from the suburbs of Chennai (Tambaram) in 1966 (Giri *et al.* 2001) c. 80 km south of Sriharikota. Though

close to Chennai, our record of the species in Sriharikota deserves attention, as the species has not been recorded from Andhra Pradesh. Additionally, the record is significant due to the insular nature of the landmass, with the Bay of Bengal to the east and Pulicat Lake on its northern, eastern and southern borders. There is also a possibility of the species occurring in other coastal areas to the north of Sriharikota in Andhra Pradesh.

A voucher specimen has been deposited in the BNHS Collection, Regn No. BNHS 4190.

May 14, 2003

S. SIVAKUMAR  
RANJIT MANAKADAN  
VARADGIRI  
Bombay Natural History Society,  
Hornbill House, S.B. Singh Road,  
Mumbai 400 023, Maharashtra, India.  
Email: bnhs@bom4.vsnl.net.in

## REFERENCES

- DANIEL, J.C. (2002): The Book of Indian Reptiles and Amphibians. Bombay Natural History Society, Mumbai. P. 181.
- DANIELS, R.J.R. (1997): A field guide to the frogs and toads of the Western Ghats, India: Part-1. *Cobra* 27: 1-25.
- DUTTA, S.K. (1997): Amphibians of India and Sri Lanka (Checklist and Bibliography). Odyssey Publishing House, Bhubaneswar. Pp. 57-58.
- GIRI, V., V. HEGDE & V. PATIL (2001): Occurrence of painted kaloula *Kaloula taprobanica* (Family Microhylidae) at Point Calimere, Tamil Nadu. *J. Bombay nat. Hist. Soc.* 98(1): 121.